



Effects of climate change on the valley glaciers of the Italian Alps

Rossana Serandrei-Barbero¹, Sandra Donnici², Stefano Zecchetto³

¹formerly ISMAR - Institute of Marine Sciences, CNR - National Research Council of Italy, Venice, Italy and Comitato Glaciologico Italiano, Torino, Italy

5 ²IGG - Institute of Geosciences and Earth Resources, CNR - National Research Council of Italy, Padova, Italy

³ISP - Institute of Polar Sciences, CNR - National Research Council of Italy, Padova, Italy and Department of Electrical Engineering, Persian Gulf University, Bushehr, Iran

Correspondence to: Sandra Donnici (sandra.donnici@cnr.it)

Abstract. The behaviour of the valley glaciers of the Italian Alps as a result of the climate changes expected for the 21st century has been investigated. From 1980 to 2017 the average length reductions of these glaciers has been 16% and their average areal reduction around 22%, much smaller than the overall glacier retreat of the Alps. Their mean observed shortening was about 500 m for a temperature increase of 1.4 °C. To quantify the valley glacier life expectancy, a model estimating their length variations from the air temperature variations of the EuroCordex climatological projections of six different models under RCP4.5 and RCP8.5 scenarios has been used. The ensemble mean temperatures in the Italian Alps region under these

15 scenarios indicate increases of temperature of ~2°C and ~4°C from 2018 to 2100 respectively. In both scenarios, the glacier model projections show a constant retreat until the eighties, weakening towards the end of the century. As expected, it resulted more severe under the RCP8.5 (from 22% to 48%) than under the RCP4.5 (from 10% to 25%) scenario, with a mean length shortening of 35% and 13% respectively by 2100. The model used estimates that the majority of the valley glaciers could better resist the climate change.

20 1 Introduction

The fluctuations of glacier fronts are mainly due to their response to year-to-year variability of the air temperature and precipitation. This variability alone can explain kilometric variations in length on a secular scale also in constant climatic conditions (Roe, 2011). The retreat rate of the last decades leaves no doubt about the incidence of the current climate anomaly and focuses interest on the behaviour of glaciers and the incidence of different morphometric factors on their shrinkage (e.g.

25 Charalampidis et al., 2018). The highly variable response of glaciers to climate changes involves changes in length, area and flow velocity that complicate the assessment of future glacier behaviour (Carturan et al., 2013). The acceleration shown by the glacial retreat in recent decades also raises new questions about the glacier life expectancy also regarding the influence of their geometry on their response to climate variations.

In recent decades, the response of glaciers to the rise in temperature since the 1980s has been analyzed using models that use the concept of climatic sensitivity (i.e., the relation between glacier length decrease and temperature increase) and attribute to





temperature the role of main forcing (Oerlemans, 2005, 2012). Models frequently use mass balance (Brown et al. 2010; Braithwaite et al. 2013; Christian et al. 2018), but a generalization of the glacier behaviour, and more specifically how the glaciers respond to the climatological variations, seems also obtainable through models dealing with the glacier length variations, given the large amount of data that are collected annually at their terminus, at present available at the World Glacier

- 35 Monitoring Service (2017 and earlier reports). For this reason, several studies have analyzed the effects of climate changes on glacier snout fluctuations (Calmanti et al., 2007; Bonanno et al., 2014; Nigrelli et al., 2015; Peano et al., 2016). The application of the model proposed by Oerlemans (2005) to the Tauern Alps glaciers (Eastern Italian Alps), obtained after a recalibration of the climate sensitivity and response time, verified its ability to reproduce the retreat of the glacier fronts even for the glaciers for which length fluctuations data are not available (Zecchetto et al., 2017). On the Tauern Alps in the last 35
- 40 years, the frontal retreat of valley glaciers has been less than that of mountain glaciers and this lower climatic sensitivity is also reflected in the projections provided by the model, which are less dramatic for the valley glaciers with respect to mountain glaciers (Serandrei-Barbero et al., 2019). The present work extends these analyses to all the valley glaciers located on the Italian side of the Alps from 2018 to 2100. It is aimed to study their future behaviour in terms of their length reduction under the increase of air temperature according to mid-range and high-end scenarios, i.e. the Representative Concentration Pathways 45 (RCPs) RCP4.5 and RCP8.5 scenarios.
 - The study is structured as follows: Section 2 introduces the glacier model used in this work. Section 3 illustrates the data employed, that is the observed data of the glaciers length and temperature and precipitation since 1980, and the climatological data obtained from Euro-Cordex (Jacob et al., 2014) (<u>https://euro-cordex.net/</u>). Section 4 presents the results, expressed in terms of the observed glaciers retreat from 1980 to 2017 and of climatological projections, then followed by Section 5 devoted to the results discussion. Section 6 summarizes the results.

2 The glacier model

50

The glaciers of the Italian Alps studied in this work are regularly observed through measurements of their annual snout position: the main parameters describing their morphological characteristics, i.e. the slope and length, are known but other important parameters such their ice thickness are unknown. This limited drastically the choice of the glacier model to use for the climatological projections: in fact, the available glacier (slope, length and snout fluctuations) and meteorological (precipitation and air temperatures) experimental data led us to address to the model proposed by Oerlemans (2005), simply relating the glacier length fluctuations L'(t) to the air temperature T'(t) fluctuations. Others more sophisticated glacier models, such as that by Roe and O'Neal (2009), cannot be applied to the valley glaciers of the Italian Alps essentially because the ice thickness is not available. The model adopted in this work writes as

$$60 \quad \frac{dL'(t)}{dt} = -\frac{C_{\rm s} T'(t) + L'(t)}{\tau} \,,$$

(1)



(4)

where C_s is the climate sensitivity (m K⁻¹) and τ is the glacier response time (year). L'(t) (m) is the variation in the glacier length with respect to its average value over the period considered. The climate sensitivity C_s is defined as

$$C_s = \frac{p_{ann}^{1/2}}{c_1 * s},$$
(2)

variations. where \bar{P}_{ann} (m y⁻¹) is the mean annual precipitation at the glacier site and s is the glacier slope. The response time 65 τ is

$$\tau = \frac{c_2}{c_3 \bar{\rho}_{ann}^{1/2} s(1+20s)^{1/2} L^{1/2}} , \qquad (3)$$

where the constant $c_3 = 0.006$ was obtained from calibration with numerical simulations (Oerlemans, 2005) and $c_1 = 0.0078 \pm 0.0004$ and $c_2 = 1.35 \pm 0.14$ result from a re-calibration carried out by Zecchetto et al. (2017) on the Italian Eastern Alps. \tilde{L} is the characteristic glacier length over the considered period.

70 Both C_s and τ depend on the morphological characteristics of the glaciers, i.e. glacier length \tilde{L} and slope s, and on the mean annual precipitation \bar{P}_{ann} . Thus, they are not independent of each other. Under similar morphological characteristics, glaciers located in more rainy locations have larger C_s and smaller τ than those in dryer areas. On the contrary, in areas of similar precipitation rates, steeper glaciers have smaller C_s and τ .

The model can be used (Oerlemans, 2011; Leclercq and Oerlemans, 2012) if

75
$$\sigma_L/\tilde{L} \ll 1$$
,

where σ_L is the standard deviation of the glacier length and \tilde{L} the characteristic glacier length over the considered period. This condition states that the model can be applied only if the glacier variations are small with respect to its length; thus, implicitly, that the the model does not account for all the non-linear and local factors influencing a glacier's life. For the climatological projections, Eq. 4 has been computed over time windows of 15 years. The results provided in this paper are all compatible

80 with Eq. 4, and all the conclusions reached must be viewed in light of this constraint. In the climatological projections spanning long periods, it is unlikely that the values of mean annual precipitation are constant with time; also the representative glacier length \tilde{L} cannot be taken as constant, since the expected glacier reduction due to climate temperature rise.

In this work, we modified the original formulation of Oerlemans (2005) reported in Eq. 1 allowing variations of \tilde{L} and \bar{P}_{ann} 85 with time, impling that C_s and τ (Eqs. 2 and 3) are not constant any more but time dependent. Therefore, Eq. 1 becomes:

$$\frac{dL'(t)}{dt} = \frac{C_S(t)T'(t) + L'(t)}{\tau(t)},$$
(5)

a first order non-linear differential equation of the type $\frac{dL'(t)}{dt} = F(t, L')$,

with
$$F(t,L') = -\frac{c_s(t)T'(t)+L'(t)}{\tau(t)}$$
, $C_s(t) = \frac{\bar{P}_{ann}(t)^{1/2}}{c_1 * s}$, and $\tau(t) = \frac{c_2}{c_3 \bar{P}_{ann}(t)^{1/2} s(1+20s)^{1/2} L(t)^{1/2}}$







where $\bar{P}_{ann}(t)$ and $\tilde{L}(t)$ are the low frequency climatological precipitation and representative glacier length computed over a 90 15 years time window. Tests run considering C_s and τ constant or slowly variable yielded differences smaller than 5% of the determination of the glacier length variations.

In the climatological projections, we have used Eq. 5, solved numerically using the well established Runge-Kutta method (Atkinson, 1989).

3 The data

95 The data presented in this section are of two kinds: the observed data, available from 1980 to 2017, which include the glaciers snout positions and the air temperatures and precipitation at annual frequency, and the climatological model data derived from the Euro-Cordex projections until 2100.



Figure 1: The position of the glaciers and in-situ climatological stations considered in this work. Circles indicate the stations providing air temperature, crosses the precipitation.

3.1 The glacier data

Given the subjectivity of the primary glacier classification and the possible transition of some glaciers to a different typology due to their severe recent decline, this study examines only the italian glaciers classified as valley glacier, i.e. glaciers flowing down a valley and in consequence having a distinct tongue (Cogley et al., 2011). Although some other Italian glaciers are







- 105 defined as valley glacier in literature (see for example Bonanno et al., 2014), those classified as valley glaciers in the Italian Glacier Inventory (Smiraglia and Diolaiuti, 2015) represent 3% of the totality, that is 25 glaciers (96.06 km²) compared to a total of 903 glaciers (369.90 km²), 57% being mountain glaciers and 40% glacierets. Valley glaciers are few, but their length variations are measured since long time; the presence of the valley tongue, generally easily accessible, and their frequent large dimensions favoured the choice of them for the annual measurements, i.e the distance year to year between some points of the
- 110 glacier terminus and the corresponding fixed landmarks on the ground. We consider in this work the 25 glaciers reported as valley glaciers by Smiraglia and Diolaiuti (2015), mapped in Fig. 1, which reports their position along with that of the in-situ meteorological stations providing air temperature and precipitation. The area including them is a part of the Alps of about 700 km in longitude and 200 km in latitude. Among them, three glaciers have been discarded from analysis: the Miage glacier is a debris-covered glacier, the front of
- 115 which is inaccessible; the Belvedere glacier between 2001 and 2005 was travelled by a surge, a phenomenon believed to be cyclical (Van de Wal and Oerlemans, 1995); the Predarossa glacier is split since 1993 into two sectors at different altitudes and scarcely connected to the sides of a rocky outcrop: an upper-altitude sector seems active as a probable circus glacier, a lower-altitude sector, extended along the slopes on the orographic left, is fed by avalanche and classifiable as slope glacier. For 19 of the remaining 22 valley glaciers, the annual ground measurements of snout fluctuations collected by the Italian
- 120 Glaciological Committee are available since the 70s of the last century, that is, from the beginning of the last positive fluctuation before the current retreat. Through the annual reports of these measurements (CGI, 1981-2018), the length and slope of the considered glaciers were updated to 2017. For four of them, snout measurements stopped before 2017 due to major morphological changes induced by the predominant processes of down-wasting during the early 21st century: on the Tza de Tzan glacier the measurements at the front stopped in 2001 with the emergence of the rocky escarpment which prevents the
- 125 feeding of the tongue and converts the valley glacier into a hanging glacier; for the Lys Glacier, snout measurements stopped in 2008 when the valley tongue separated from the feeder basin; on the Pisgana glacier, since 2011 the tongue has been interrupted by a rocky outcrop that limits its feeding; finally, the Verra Grande glacier tongue is not accessible since 2014 for the repositioning of the glacier terminus above a rocky outcrop. The length variations of the three glaciers without measurements of their terminus fluctuations (Osand Sud, Vitelli and Vallelunga glaciers) have been obtained from the historic
- 130 observed temperature anomalies through the model illustrated in Section 2. Table 1 shows the main morphometric parameters of the 22 valley glaciers considered. The glacier length and area data were collected on the ground between the end of the 70s and the early 80s during the campaigns for the compilation of the World Glacier Inventory (WGMS, 2017 and earlier reports) and 1980 constitutes the reference year for the length and area values. The lengths in 2017 are from the annual field measurements of Comitato Glaciologico Italiano (1981-2018) starting from 1980
- 135 lengths. The 2015 area values are derived from orthophotos or satellite images taken between 2007 and 2011 (Smiraglia and Diolaiuti, 2015).







Between 1980 and 2017, the 22 considered valley glaciers lost an average length of 542 m and an average area of 0.88 km² for a total loss of 19.36 km². In the same period on the Italian side of the Alps the glacierized total area loss was 238.66 km² corresponding to 39% with respect to the 1980 glacierized total area (Smiraglia and Diolaiuti, 2015).

140

Table 1: Main parameters of the 22 valley glaciers considered. Slope is the angle between the glacier surface and the horizon. The	e
values of climate sensitivity C_s and response time τ were computed by Eqs. 2 and 3.	

Alpine sector	Glacier	WGI code	latitude N	longitude E	Area 1980 (km ²)	Area 2015 (km ²)	∆ area (%)	Length 1980 (m)	Length 2017 (m)	∆ length (%)	Slope 1980 (°)	Cs (mK ⁻¹)	τ (years)
Western Alps	Soches-Tsanteleina	IT4L01514011	45° 28' 51"	7° 03' 41"	3.32	2.77	-17	3400	2781	-18	12	571.6	7.1
	Gliairetta-Vaudet	IT4L01515021	45° 30' 17"	7° 01' 31"	4.41	3.72	-16	3200	2810	-12	19	333.0	3.3
	Tza de Tzan	IT4L01522024	45° 58' 40"	7° 33' 44"	3.95	3.27	-17	3700	3600 (2001*)		19	387.3	3.9
	Verra Grande	IT4L01504004	45° 55' 34"	7° 45' 33"	7.28	6.6	-9	5100	4425 (2014*)		18	421.8	3.6
	Lys	IT4L01502002	45° 54' 31"	7° 49' 58"	11.82	9.89	-16	5600	5357 (2008*)		20	319.1	2.4
	Osand Sud	IT4L01216015	46° 23' 34"	8° 19' 39"	3.64	2.21	-39	2810	1712**	-39**	15	491.1	6.1
	Osand Nord	IT4L01216018	46° 24' 29"	8° 18' 20"	1.98	1.31	-34	2878	2703	-6	14	618.6	8.5
	Ventina	IT4L01122009	46° 16' 02"	9° 46' 17"	2.26	1.89	-16	3171	2684	-15	23	291.5	3.1
	Vitelli	IT4L01139003	46° 30' 03"	10° 27' 39"	1.82	1.89	4	3100	2174**	-30**	16	442.7	5.5
	Cedech	IT4L01137018	46° 27' 00"	10° 36' 22"	2.84	2.07	-27	3085	2565	-17	20	304.5	3.4
	Forni	IT4L01137024	46° 23' 32"	10° 35' 28"	13.24	11.34	-14	5062	4363	-14	15	449.5	4.7
	Dosegù	IT4L01137031	46° 22' 34"	10° 33' 07"	3.44	2.16	-37	3149	2606	-17	14	521.8	7.2
	Pisgana Ovest	IT4L01028006	46° 11' 59"	10° 32' 40"	2.47	2.49	1	2650	2190 (2010*)		15	415.5	5.8
	Venerocolo	IT4L01028011	46° 09' 46"	10° 29' 56"	1.18	0.82	-31	2144	1968	-8	17	544.9	9.2
Eastern Alps	Presanella	IT4L00102413	46° 13' 35"	10° 39' 24"	3.92	2.79	-29	3200	3093 (2007)		18	330.9	3.8
	La Mare	IT4L00102516	46° 26' 05"	10° 38' 00"	4.75	3.53	-26	3470	2982	-14	24	337.3	3.6
	Forcola	IT4L00112125	46° 27' 07"	10° 38' 26"	2.52	1.77	-30	3500	2596	-26	19	361.0	4.0
	Cevedale	IT4L00112126	46° 27' 20"	10° 37' 45"	3.20			3700	2841	-23	17	411.7	4.7
	Vedretta Lunga	IT4L00112128	46° 28' 05"	10° 36' 55"	3.22	1.76	-45	3600	2680	-26	13	537.0	7.0
	Solda	IT4L00112417	46° 29' 23"	10° 34' 05"	6.48	5.5	-15	4200	4020	-4	21	346.7	3.5
	Vallelunga	IT4L00112907	46° 48' 41"	10° 43' 57"	8.55	7.35	-14	3900	3180**	-18**	20	344.7	3.5
	Gran Pilastro	IT4L00121313	46° 57' 51"	11° 43' 45"	2.62	1.73	-34	3700	3060	-17	14	402.2	6.5
	mean values				4.54	3.66	-22	3560	3018	-16	17	417.5	5.0

* glaciers not considered in the projections ** unmeasured glacier

3.2 The meteorological data

- 145 The air temperature and total precipitation used in this work are both historical and forecast data. The former, obtained from the HISTorical instrumental climatology surface time series of the greater ALPine region (HISTALP, www.zamg.ac.at/histalp/, Auer et al., 2007), are available from 1980 to 2014. The stations selected are positioned between 1300 m and 2470 m of altitude: their location is reported in Fig. 1, with circles and crosses.
- Figure 2 reports the historical time series of temperature fluctuations (left panel) and annual total precipitation (right panel) 150 derived from the data of the available stations; the latter sorted for their longitudinal location. The temperature fluctuations, i.e. the temperature values with respect to their mean, exhibit a similar trend, an increase of about 1.4° C in the period 1980-2015, while the total precipitation show no trends but larger values in the western than in the eastern part of the region under study. Total precipitation data have been used to compute C_s and τ (see Section 2, Eqs. 2 and 3) for all the glaciers. The air



155

170



temperature variations have been used instead to reproduce the length historic variations of the three glaciers not measured (Osand Sud, Vitelli and Vallelunga glaciers).

The atmospheric climatological model data used are the Euro-Cordex (Jacob et al., 2014) (https://euro-cordex.net/) temperature



Figure 2: Time series of experimental air temperature variations (left panel) and annual total precipitation (right panel) from the in-situ stations available.

and precipitation projections of the climatological Representative Concentration Pathways (RCPs) RCP4.5 and RCP8.5 scenarios. The Euro-Cordex are regional climate model simulations for the European domain with resolution of 0.11 degree (EUR-11, ~12.5km). The two climatological scenarios are based on greenhouse gas emission corresponding to stabilization of

- 160 radiative forcing after the 21st century at 4.5 W/m² (RCP4.5), rising radiative forcing crossing 8.5 W/m² at the end of 21st century (RCP8.5) (Moss et al., 2010; Nakićenović et al., 2000; Van Vuuren et al., 2008). Six different regional climatological models providing air temperature and total precipitation, i.e. CCLM4-8-17.v1 (Rockel et al., 2008), HIRHAM5.v3 (Christensen et al., 1998), Racmo2.2 (Van Meijgaard et al., 2012), RCA4.v1 (Samuelsson et al., 2011), WRF (Skamarock et al., 2008), and REMO2009 (Jacob et al., 2012), have been used in order to derive ensemble averages of the glacier future
- 165 length variations. In the following, we will refer as ensemble averages the mean of the time series from the different climatological models.

The ensemble air temperature anomalies at the glaciers position are reported in Fig. 3 for the two scenarios considered. While for the milder RCP4.5 there will be a rise of temperature of about 1.6° C from 2020 to 2086 and then a light decrease to 2100, RCP8.5 indicates an increase of about 4.5° C from 2020 to 2095, followed by a light decrease of some tenths of a degree. The mean increase of temperature is $0.020 \pm 0.014^{\circ}$ C/y and $0.053 \pm 0.017^{\circ}$ C/y respectively.

In some period, the air temperature variations T' vary significantly (~ 0.3 °C) according to the glacier size, as in the period 2020-2040 and 2080-2100. This is because the simulations have been carried out using the climatological temperature and precipitation closest to the glacier location.







4 Results

175 4.1 Glaciers retreat until 2017

Between 1980 and 2017 the loss of length of the valley glaciers was between 4% and 26% with an average contraction of 16% (about 500 m). The areal reduction was between 9% and 45% with an average shrinkage around 22%. Table 1 provides an overview of the length and area variations that occurred between 1980 and 2017 together with the C_s and τ values obtained from Eqs. 2 and 3. The climate sensitivity C_s ranges between 291 m K⁻¹ (Ventina Glacier) and 619 m K⁻¹ (Osand Nord Glacier)

180 with a mean value of 417 m K⁻¹. A temperature increase of 1.4 °C from 1980 to 2015 produces a mean length decrease of 584 m, consistent with the observed average shortening of 542 m. The glacier response time τ is between 2 and 10 years, with a mean value of 5 years.



Figure 3: The ensemble air temperatures variations over the six regional models at the grid points closest the glaciers position from 2018 to 2100. Left panel: RCP4.5 scenario. Right panel: RCP8.5 scenario. The colours are sorted from the longest (blue) to the shortest (red) glacier.

The climate sensitivity C_s (Eq. 2) depends on the glacier slope and the total annual precipitation. Glaciers with slope $18^\circ < s < 27^\circ$ have $C_s < 350$ m K⁻¹ and response times $1.7 < \tau < 3.8$ years; glaciers with a more gentle slope $12^\circ < s < 16^\circ$ have $C_s > 450$ m K⁻¹ and τ of 7 - 9 years. Figure 4 reports *Cs* versus the glacier slope *s* for the 22 valley glaciers considered in this work: as 190 expected by Eq. 2, *Cs* decreases, as s increases, in average – 24 m for one degree of slope.







Figure 4: C_s versus the glaciers slope s for the 22 valley glaciers considered in this work. The solid line is the linear best fit. The glaciers in the legend are sorted from the longest (blue) to the shortest (red).





Figure 5 reports the observed retreat from 1980 to 2017 of the measured glaciers: all exhibit a continuous shrinkage since
1990, but with very different rates, from ~7% of the Solda and Osand Nord glaciers (respectively among the longest and the shortest) to ~25% of Forni and Forcola glaciers.

4.2 Glaciers projections to 2100

Climatological glacier lengths have been obtained forcing the glacier model (Eq. 5) with the annual air temperature variations and the total precipitation derived by each of the six atmospheric climatological models chosen (see Section 3.2) at the grid

200 point closest to the glacier location.

The results are shown as ensemble means, derived averaging the glacier lengths variations obtained from each atmospheric climatological model. The Tza de Tzan, Verra Grande, Lys, and Pisgana W glaciers, that before 2017 underwent morphological variations, cannot be anymore considered valley glaciers. Therefore, they were not considered in the projections, which include instead the three unmeasured glaciers (Osand Sud, Vitelli and Vallelunga), the 2017 length of which derived by the glacier

205 model simulations. Consequently, the climatological projections have been carried out over 18 glaciers. Figure 6 reports the ensemble glacier length variations in percent from 2018 to 2100 under the RCP4.5 (top panel) and RCP8.5 (bottom panel) scenarios. These have been obtained averaging the projections derived by every single climatological model



Figure 6: The percent glaciers length variations derived from climatological projections from 2018 to 2100. Top panel: RCP4.5 scenario. Bottom panel: RCP8.5 scenario. The colours are sorted from the longest (blue) to the shortest (red) glacier.





respecting the glacier model constraint of Eq. 4, computed over time windows of 15 years, set as $\sigma_L/\tilde{L} < 0.1$, i.e. glacier variations at least one order of magnitude smaller than the glacier length.

Figure 7 shows the ensemble values of σ_L/\tilde{L} : for RCP4.5 scenario (top panel) $\sigma_L/\tilde{L} < 0.05$, thus respecting the constraints of Eq. 4 for the whole projections period; for scenario RCP8.5 (bottom panel) σ_L/\tilde{L} is greater than 0.1 in the late seventies for two glaciers (Dosegu and Venerocolo), which therefore cannot be projected until 2100. These glaciers will probably disappear within the end of the century.

In both scenarios, there is a constant retreat until the eighties of this century, weakening towards the end of the century. As

215 expected, it is more severe under the RCP8.5 (from 22% to 48%) than under the RCP4.5 (from 10% to 25%), since the almost double temperature rise foreseen by the farther scenario (see Fig. 3).







Figure 7: Ensemble values of σ_L/\tilde{L} derived from climatological projections from 2018 to 2100. Top panel: RCP4.5 scenario. Bottom panel: RCP8.5 scenario. The colours are sorted from the longest (blue) to the shortest (red) glacier.

220 Figure 8 shows the mean percent of the glacier length reductions shown in Fig. 6: by 2100 there will be a loss of 35% and 13% of the glacier lengths under RCP8.5 and RCP4.5 respectively. The error bars, accounting both for the different behaviour of the glaciers under the same climatological forcing and for the different meteorological forcing provided by the six different





climatological model used, are always smaller than \pm 7%, making reliable the figure of the mean retreat. Therefore the different climatological models impact on the glacier model results for less than 10%.



225

Figure 8: The mean percent glacier length reduction from 2018 to 2100 according to the RCP4.5 and RCP8.5 climatological scenarios.

These results indicate that, even under the most severe RCP8.5 scenario, the majority of glacier length variations will not be so large as to menace the existence of the glaciers to 2100 with the exceptions of two (Venerocolo and Dosegu), which will

230 preserve the present behaviour characterized by glacier size much bigger than their annual variations. This even if some of them, i.e. Vitelli, Osand Nord and Vedretta Lunga glaciers, will almost half their 2018 length, in accordance with the general trend that attributes greater sensitivity to gently sloping glaciers (Fig. 4).





5 Discussion

Studies dealing with alpine glaciers generally group them based on their geographic location (Carturan et al., 2013; Fischer et
al., 2015 among others) or their size (DeBeer and Sharp, 2009; Huss and Fischer, 2016) while their primary classification is not considered, which seems to have some influence on the climatic sensitivity.
On the valley glaciers, from 1980 to 2017 the mean length loss is 16%. Their average areal shrinkage of 22% between 1980

and 2015 (Table 1) shows their smaller retreat with respect to the general shrinkage in the European Alps estimated around 40% as a lower bound (Knoll and Kerschner, 2009; Lambrecht and Kuhn, 2007; Nigrelli et al., 2015; Paul, 2002; Zemp et al.,

240 2006, 2015).

260

On the Eastern Alps, a previous comparison between valley and mountain glaciers belonging to the same size classes highlighted the lower retreat of valley glaciers (Serandrei-Barbero et al. 2019).

All the 22 valley glaciers considered in this paper have an area > 1 km², except one, while 90% of the glaciers on the Italian Alps (812 glaciers) have areas < 1 km². In general larger glaciers show regresses inversely proportional to their size (Serandrei-

- Barbero et al. 1999; Paul et al. 2004, 2011; Diolaiuti et al. 2012; Carturan et al. 2013); this could significantly contribute to their expected minor retreat. However, on the valley glaciers considered here it does not seem to be any relationship between size and frontal retreat as it can be inferred from Fig. 5, which reports past percent retreats sorted according to the glacier size. The frontal variations, as reported in Section 4.1, result influenced by the slope, in accordance with Hoelzle et al. (2003) and Huss and Fischer (2016), the latter describing values of Cs and τ generally higher on gently sloping glaciers. The longer
- 250 response time of gently sloping glaciers, slowing down the feeding of the valley tongue, would favour processes of nonlinear decay. On the contrary, the steepness of the glacial mass, which is accompanied by a shorter response time, seems to favour the feeding of the tongue, leaving rise to lower Cs and minor withdrawals. The response time τ obtained, between 2 and 10 years, agree with τ values of 2-4 years of the Kesselwandferner glacier and the Palù glacier on the northern side of the Eastern Alps (Oerlemans 2007) and, more generally, with τ values of 2-21 years of the glaciers of the Southern side of the Eastern
- Alps (Serandrei-Barbero et al., 2019) of length comparable with the glaciers considered here. However, the response times of valley glaciers are small compared to the response times between 10 and 20 years generally observed in the Alps (Paul et al. 2004).

The results of the glaciers shrinkage due to the simulated changes of air temperature and total precipitation as foreseen by six atmospheric models under the RCP4.5 and RCP8.5 climatological scenarios are slightly modulated (2%) by the variations of the climatological total precipitation which enters in the definition of the glacier sensitivity Cs and response time τ .

The main question concerns the reliability of these estimates. A partial answer to this crucial issue may be derived looking to the observed retreats shown in Fig. 5 and comparing them with those from the two climatological scenarios in terms of retreat velocity. The mean retreat velocity from 1980 to 2017 was 15±7 m/y (with a mean length loss of about 500 m), while that of the climatological scenarios RCP4.5 and RCP8.5, computed over period of the same temperature increase occurred from 1980





265 to 2017 (~1.4°C), is 10±3 m/y and 15±5 m/y respectively. The glacier retreat speed under the RCP8.5 scenario is very close to that observed in the period 1980-2017.

As for the observed glacier length retreats (Fig. 5), the climatological results reported in Fig. 6 show that glaciers behave differently under the same climatological forcing: under the RCP4.5 scenario, there is up to 20% of difference between the glaciers with smaller reduction and those of largest percent retreat, which almost doubles under RCP8.5. These differences are

- 270 possibly due to the values of Cs and τ of each glacier and, in consequence, to their characteristic length and slope, as well as to the annual total precipitation in the glacier site. But other variables could influence the glacier behaviour mainly due to differences in the geometric characteristics and topographic features of individual glaciers (Paul et al. 2004; Haeberli et al. 2007; Oerlemans 2012; Huss and Fischer 2016 among others). Furthermore, the model cannot take into account morphometric variations such as changes in the elevation range related to frontal retreat or variations in glacier hypsometry (Kuhn 1985; Paul
- 275 and Haeberli 2008; Fischer et al. 2015; Zemp et al. 2015; Charalampidis et al. 2018). The results of the model should be considered as first order estimates, even because it cannot account for the changes introduced by positive feedbacks, such as the decrease in albedo and the thermal emission from increasing rocky outcrops, as well as possible negative feedbacks, such as the contribution of avalanches or the increasing debris cover or shaded area. However, since the majority of the glacier simulated projections satisfies the constraint that the glacier variations are much smaller than the glacier length (Eq.4), thus
- 280 longer preserving the present glacier morphology, the results of model can be considered reliable: for the valley glaciers the average length loss expected for 2100 under the most severe scenario RCP8.5 is between 13% and 60%, with a mean value of 35%. The projections obtained under RCP4.5, more favourable, show instead length reduction between 5% and 23%, with a mean value of 13%.
- According to the estimate based on the scaling relationship between glacier area and length derived for the mid-latitudes glacier regions (Machguth and Huss 2014), the observed reduction in length of 16% between 1980 and 2017, corresponds to an area loss of 22%. A rough estimate based on the same proportion between length and area decrease indicates for 2100 an area loss of about 50% (1.83 km²) for an average length loss of 35% under RCP8.5 scenario; under the RCP4.5 scenario, a length retreat of 13% corresponds to an area loss of 20% (0.73 km²).
- On the glaciers of the European Alps, the projections for the end of the century indicate volume losses between 75% and 89% under RCP4.5 and between 90% and 98% under RCP8.5 (Marzeion et al. 2012; Radic et al. 2014; Huss and Hock 2015; Zekollary et al. 2019) with the possible disappearance between 69% to 92% of all the glaciers in the Alps (Zebre et al. 2020). Both mountain and valley glaciers are included in this severe estimate, but the behaviour of the few valley glaciers is undetectable from that of the great majority of alpine glaciers.
- A comparison is instead possible with the projections of some glaciers on the southern side of the Alps: on the Italian Western Alps, on 14 glaciers considered as valley glaciers by Bonanno et al. (2014), the mean length loss projected to 2050 is quantified in 300-400 m compared to the 2010 length with the RCP4.5 scenario, with larger retreats under RCP8.5 scenario Even though these glaciers are smaller (mean area 2.79 km²) than the glaciers considered in the present study, their projected retreat is consistent with the average retreat of 7% (about 200 m) and 13% (about 400 m), expected for valley glaciers in 2050 under
 - 15





RCP4.5 and RCP8.5 respectively (Fig. 8). Moreover, based on A1B scenario (Nakicenovic et al. 2000), also on the Eastern
Alps the expected length loss in 2100 between 20% and 35% on the valley glaciers is much less than the mean shortening between 35% and 60% expected on mountain glaciers of the same size class (Serandrei-Barbero et al. 2019). The smaller decline of the valley glaciers with respect to the totality of Alpine glaciers is also confirmed by the comparison with the projections of Peano et al. (2016): on the studied mountain glaciers in the Italian Western Alps the expected average length shrinkage in 2100 under the future scenarios RCP4.5 and RCP8.5 is respectively 47% and 68.5% and the length would become zero in 2100 for some of these mountain glaciers.

- On both valley glaciers and mountain glaciers, length reductions are expected also in case of an attenuation or inversion of the warming in progress because of the response time τ , due to the imbalance always present between the transient length and the length in conditions of reached equilibrium (Christian et al. 2018). Citterio et al. (2007) suggest that, in the coming decades, only the few largest glaciers will be in a condition to survive. The last 35 years glacier behaviour shown in the present study
- 310 indicates a minor sensitivity of the valley glaciers compared both to mountain glaciers of Eastern Alps, analyzed with the same model (Serandrei-Barbero et al., 2019), and to the generality of alpine glaciers, as reported in literature. Therefore, the glacier size and typology could both play a decisive role in the behaviour of valley glaciers under the climate change, but their projections obtained from the model suggest, for valley glaciers, a possible less dramatic trend than expected.

315 6 Conclusions

Although the literature indicates that the majority of glaciers will disappear in the coming decades, the valley glaciers seems able to better resist the changing climate and the majority of them could be still present up the end of 21st century. Between 1980 and 2017 the valley glaciers of the Italian Alps lost on average 16% of their length and 22% of the surface, a much lower value than the general retreat of the glaciers of the Alps. By 2100, the model projections under the RCP4.5 and

- 320 RCP 8.5 climatological scenarios indicate for these glaciers a mean shortening of 13% and 35% respectively, a much lower value than the expected retreat of the majority of glaciers in the Alps, as reported in literature. The differences of observed length reductions are up to 20% in the last 40 years. In the projections, for the next 40 years, the differences are up to 18% for the scenario RCP8.5 (Fig. 6), consistent with those observed. Summarizing our results: a) under the RPC8.5 scenario, the mean retreat velocity of the glaciers is similar to that of the period
- 325 1980 to 2017 obtained from the observed data; b) given the form of the model used, the main forcing is the air temperature variation, while total precipitation changes have a weak impact on the results; c) the impact of different climatological models on the simulated results is less than 10%; d) according to the model, under the RCP8.5 scenario the majority of the valley glaciers (about 80%) will resist the climate change experiencing retreats less than 50% of their 2017 length and thus probably maintaining their characteristics of valley glaciers.





- 330 The simulated glacier retreats have been derived by a glacier model that does not account for possible non-linear effects and consequent geometric changes. Furthermore, possible disintegration and down wasting could deliver some of the valley glaciers to the typology of mountain glaciers, threatened by a much more pronounced retreat. Moreover, progressive glacier shrinkage and fragmentation will lead to a general increasing glacier melt even under the same climatic conditions. Despite the effect of climate change indicates on the Alps a severe ice masses decline, due to the minor retreat of the valley
- 335 glaciers in the last 35 years with respect to the generality of the alpine glaciers, the model suggests that the valley glaciers are less sensitive to the air temperature and precipitation. Their limited length losses result in a slow glacier retreat, thus longer preserving the glacier typology and making the valley glaciers probably better resist the climate change.

Conflicts of interest/Competing interests (include appropriate disclosures): The authors declare that they have no known

340 competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. Code availability (software application or custom code) Not applicable Authors' contributions: R.S.B and. S.D. collected and analyzed the glaciers data, interpreted and discussed the results and contributed to the paper final version; S.Z. collected and analyzed the climatological data, developed the glaciers model and contributed to the paper final version.

345 Acknowledgements

The field measurements from 1980 to 2017 were supported by the Comitato Glaciologico Italiano: thanks are due to the voluntary operators for annually monitoring glacier snout fluctuations. Authors are grateful to Giovanni Mortara and Stefano Perona (Comitato Glaciologico Italiano), Michela Munari (Provincia Autonoma di Bolzano-Ufficio Idrografico) and Franco Secchieri for providing them with the original data collected on the ground during the campaigns for the compilation of the

350 World Glacier Inventory. Thanks to Marco Zecchetto (Instituto Superior Técnico, Universidade de Lisboa, Portugal) for the suggestions provided to solve the model equation. Euro-Cordex climatological data have been downloaded from esgfdata.dkrz.de/projects/cordex-dkrz.

References

Atkinson, K. E.: An Introduction to Numerical Analysis (2nd ed.), John Willey & Sons, New York, 604 pp., 1989.

355 Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C., Briffa, K., Jones, P.D., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin, J. M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S., Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Nieplova, E.: HISTALP – Historical instrumental climatological surface time series of the greater Alpine region 1760-2003, Int J Climatol, 27, 17-46, 2007.







- Bonanno, R., Ronchi, C., Cagnazzi, B., and Provenzale, A.: Glacier response to current climate change and future scenarios in the north-western Italian Alps, Reg Environ Change, 14(2), 633-643, https://doi.org/10.1007/s10113-013-0523-6, 2014.
 Braithwaite, R. J., Raper, S. C., and Candela, R.: Recent changes (1991-2010) in glacier mass balance and air temperature in the European Alps, Ann Glaciol, 54(63), 139-146, https://doi.org/10.3189/2013AoG63A285, 2013.
 Brown, J., Harper, J., and Humphrey, N.: Cirque glacier sensitivity to 21st century warming: Sperry Glacier, Rocky Mountains,
- 365 USA, Global Planet Change, 74(2), 91-98, 2010. Calmanti, S., Motta, L., Turco, M, and Provenzale, A.: Impact of climate variability on Alpine glaciers in northwestern Italy, Int J Climatol, 27, 2041-2053, https://doi.org/10.1002/joc.1500, 2007. Carturan, L., Filippi, R., Seppi, R., Gabrielli, P., Notarnicola, C., Bertoldi, L., Paul, F., Rastner, P., Cazorzi, F., Dinale, R., and Dalla Fontana, G.: Area and volume loss of the glaciers in the Ortles-Cevedale group (Eastern Italian Alps): controls and
- 370 imbalance of the remaining glaciers, The Cryosphere, 7, 1339-1359, https://doi.org/10.5194/tc-7-1339-2013, 2013. Charalampidis, C., Fischer, A., Kuhn, M., Lambrecht, A., Mayer, C., Thomaidis, K., and Weber, M.: Mass-Budget Anomalies and Geometry Signals of Three Austrian Glaciers, Front Earth Sci, 6, https://doi.org/10.3389/feart.2018.00218, 2018. Christensen, O. B., Christensen, J. H., Machenhauer, B., and Botzet, M.: Very high-resolution regional climate simulations over Scandinavia— Present climate, J Climate, 11, 3204-3229, 1998.
- Christian, J., Koutnik, M., and Roe, G.: Committed retreat: controls on gla,ier disequilibrium in a warming climate, J Glaciol, 64, 675-688, 2018.
 Citterio, M., Diolaiuti, G., Smiraglia, C., D'Agata, C., Carnielli, T., Stella, G., and Siletto, G. B.: The fluctuations of italian glaciers during the last century: a contribution to knowledge about alpine glacier changes. Geogr Ann A, 89(3), 166-184, https://doi.org/10.1111/j.1468-0459.2007.00316.x, 2007.
- Cogley, J., Hock, R., Rasmussen, L., Arendt, A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G., Müller, M., Nicholson, L., and Zemp, M.: Glossary of glacier mass balance and related terms, IHP-VII Technical Documents in Hydrology No. 86, IACS Contribution No. 2, UNESCO-IHP, Paris, 2011.
 CGI Comitato Glaciologico Italiano: Relazioni delle campagne glaciologiche 1980-2017, Geogr Fis Din Quat, 4-41, 1981-2018.
- 385 DeBeer, C. M., Sharp, M.: Topographic influences on recent changes of very small glaciers in the Monashee Mountains, British Columbia, Canada, J Glaciol, 55, 691-700, 2009. Diolaiuti, G. A., Bocchiola, D., Vagliasindi, M., D'Agata, C, and Smiraglia, C.: The 1975-2005 glacier changes in Aosta Valley (Italy) and the relations with climate evolution, Prog Phys Geog, 36(6), 764-785, 2012. Fischer, M., Huss, M., and Hölzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980-2010, The Cryosphere,
- 9(2), 525-540, https://doi.org/10.3929/ethz-b-000099894, 2015.
 Haeberli, W., Hoelzle, M., Paul, F., and Zemp, M.: Integrated monitoring of mountain glaciers as key indicators of global climate change: the European Alps, Ann Glaciol, 46, 150-160, https://doi.org/10.3189/172756407782871512, 2007.





Hoelzle, M., Haeberli, W., Dischl, M., and Peschke, W.: Secular glacier mass balances derived from cumulative glacier length changes, Global Planet Change, 36(4), 295-306, 2003.

395 Huss, M., Fischer, M.: Sensitivity of very small glaciers in the Swiss Alps to future climate change, Front Earth Sci, 4(34), 1-17, https://doi.org/10.3389/feart.2016.00034, 2016.

Huss, M., Hock, R.: A new model for global glacier change and sea-level rise, Front Earth Sci, 3, 1-22, https://doi.org/10.3389/feart.2015.00054, 2015.

Jacob, D., Elizalde, A., Haensler, A., Hagemann, S., Kumar, P., Podzun, R., Rechid, D., Remedio, A. R., Saeed, F., Sieck, K.,

400 Teichmann, C., and Wilhelm, C.: Assessing the transferability of the regional climate model REMO to different coordinated regional climate downscaling experiment (CORDEX) regions, Atmosphere-Basel, 3, 181-199, https://doi.org/10.3390/atmos3010181, 2012.

Jacob, D., Petersen, J., Eggert, B., Alias, A., Bössing Christensen, O., Bouwer, L. M., Braun, A., Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K.,

- 405 Kovats, S., Kröner, N., Kotlarski, S., Kriegsmann, A., Martin, E., van Meijgaard, E., Moseley, C., Pfeifer, S., Preuschmann, S., Radermacher, C., Radtke, K., Rechid, D., Rounsevell, M., Samuelsson, P., Somot, S., Soussana, J., Teichmann, C., Valentini, R., Vautard, R., Weber, B., and Yiou, P.: EURO-CORDEX: new high-resolution climate change projections for European impact research, Reg Environ Change, 14, 563-578, https://doi.org/10.1007/s10113-013-0499-2, 2014. Kuhn, M., Markl, G., Kaser, G., Nickus, U., Obleitner, F., and Schneider H.: Fluctuations of climate and mass balance: different
- responses of two adjacent glaciers, Z Gletscherk Glazialgeol, 21, 409-416, 1985.
 Knoll, C., Kerschner, H.: A glacier inventory for South Tyrol, Italy, based on airborne laser-scanner data, Ann Glaciol, 50 (53), 46-52. https://doi.org/10.3189/172756410790595903, 2009.
 Lambrecht, A., Kuhn, M.: Glacier changes in the Austrian Alps during the last three decades, derived from the new Austrian glacier inventory, Ann Glaciol, 46, 177-184. https://doi.org/10.3189/172756407782871341, 2007.
- Leclercq, P. W., Oerlemans, J.: Global and hemispheric temperature reconstruction from glacier length fluctuations, Clim Dynam, 38(5-6), 1065-1079, https://doi.org/10.1007/s00382-011-1145-7, 2012.
 Machguth, H., Huss, M.: The length of the world's glaciers-a new approach for the global calculation of center lines, The Cryosphere, 8(5), 1741-1755, https://doi.org/10.5194/tc-8-1741-2014, 2014.
 Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sealevel change from the surface mass balance of glaciers, The
- Cryosphere, 6, 1295-1322, https://doi.org/10.5194/tc-6-1295-2012, 2012.
 Moss, R. H., Edmonds, J. A., Hibbard, K., Manning, M., Rose, S. K., Van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., and Kram, T.: The next generation of scenarios for climate change research and assessment, Nature, 463, 747-756, https://doi.org/10.1038/nature08823, 2010.

Nakićenović, N., Alcamo, J., Davis, G., de Vries, B., Fennhann, J., Gaffin, S., Gregory, K., Grübler, A., Jung. T. Y., Kram,
T., La Rovere, E. L., Micaelis, L., Mori, S., Morita, T., Pepper, W., Pitcher, H., Pricer, L., Raihi, K., Roehrl, A., Rogner, H.
H., Sankovski, A., Schlesinger, M., Schukla, P., Smith, S., Swath, R., van Rooigen, S., Victor, N., and Dadi, Z.: Emissions





Scenarios. A special report of working group III of the inter-governamental panel on climate change, Technical report, IPCC, 2000.

 Nigrelli, G., Lucchesi, S., Bertotto, S., Fioraso, G., and Chiarle, M.: Climate variability and Alpine glaciers evolution in
 Northwestern Italy from the Little Ice Age to the 2010s, Theor Appl Climatol, 122(3-4), 595-608, https://doi.org/10.1007/s00704-014-1313-x, 2015.

Oerlemans, J.: Extracting a climate signal from 169 glacier records, Science, 308(675), https://doi.org/10.1126/science.1107046, 2005.

- Oerlemans, J.: Estimating response times of Vadret da Morteratsch, Vadret da Palü, Bricksdalsbreen and Nigardsbreen from 435 their length records, J Glaciol, 53, 357-362, 2007.
- Oerlemans, J.: Minimal Glacier Models. Utrecht University, Utrecht, ISBN 78-90-6701-022-1, 2011. Oerlemans, J.: Linear modelling of glacier length fluctuations, Geogr Ann A, 94(2), 183-194, 2012. Paul, F.: Changes in glacier area in Tyrol, Austria, between 1969 and 1992 derived from Landsat 5 Thematic Mapper and Austrian Glacier Inventory data, Int J Remote Sens, 23(4), 787-799, 2002.
- Paul, F., Frey, H., and Le Bris, R.: A new glacier inventory for the European Alps from Landsat TM scenes of 2003: challenges and results, Ann Glaciol, 52, 144-152, https://doi.org/10.3189/172756411799096295, 2011.
 Paul, F., Haeberli, W.: Spatial variability of glacier elevation changes in the Swiss Alps obtained from two digital elevation models, Geophys Res Lett, 35(21), https://doi.org/10.1029/2008GL034718, 2008.
 Paul, F., Kääb, A., Maisch, M., Kellenberger, T., and Haeberli, W.: Rapid disintegration of Alpine glaciers observed with
- satellite data, Geophys Res Lett, 31(21), L21402, 1-4. https://doi.org/10.1029/2004GL020816, 2004.
 Peano, D., Chiarle, M., and von Hardenberg, J.: A minimal model approach for glacier length modeling in the Western Italian Alps, Geogr Fis Din Quat, 39(1), 69-82, https://doi.org/10.4461/GFDQ. 2016.39. 7, 2016.
 Radic, V., Bliss, A., Beedlow, A. C., Hock, R., Miles, E., and Cogley, J. G.: Regional and global projections of twenty-first century glacier mass changes in response to climate scenarios from global climate models, Clim Dynam, 42, 37-58,
- https://doi.org/10.1007/s00382-013-1719-7, 2014.
 Rockel, B., Will, A., and Hense, A.: Special issue regional climatemodelling with COSMO-CLM (CCLM), Meteorol Z, 17, 347-348, https://doi.org/10.1127/0941-2948/2008/0309, 2008.
 Roe, G. H.: What do glaciers tell us about climate variability and climate change? J Glaciol, 57(203), 567-578, 2011.
- Roe, G. H., O'Neal, M. A.: The response of glaciers to intrinsic climate variability: observations and models of late-Holocene variations in the Oacific Northwest, J Glaciol, 55(193), 839-854, 2009.
 - Samuelsson, P., Jones, C., Willén, U., Ullerstig, A., Gollvik, S., Hansson, U., Jansson, C., Kjellström, E., Nikulin, G., and Wyser, K.: The Rossby Centre Regional Climate Model RCA3: model description and performance, Tellus A, 63, 4-23, https://doi.org/10.1111/j.1600-0870.2010, 2011.

Serandrei-Barbero, R., Rabagliati, R., Binaghi, E., and Rampini, A.: Glacial retreat in the 1980s in the Breonie, Aurine and 460 Pusteresi groups (eastern Alps, Italy) in Landsat TM images, Hydrolog Sci J, 44, 279-296, 1999.



465



Serandrei-Barbero, R., Donnici, S., and Zecchetto, S.: Projected effects of temperature changes on the Italian Western Tauri glaciers (Eastern Alps), J Glaciol, 1-10, https://doi.org/10.1017/jog.2019.7, 2019.

Skamarock, W.C., Klemp, J.B., Dudhia, J., Gill, D., Barker, D. M., Duda, M. G., Huang, X., Wang, W., and Powers, J. G.: A description of the advanced research WRF version 3, Technical Report NCAR/TN-475+STR, National Center for Atmospheric Research, Boulder, Colorado, USA, 2008.

Smiraglia, C., Diolaiuti, G. (eds.): The new Italian glacier inventory, Ev-K2-CNR, Bergamo Pubbl., Bergamo, 2015.
 Van de Wal, R. S. W., Oerlemans, J.: Response of valley glaciers to climate change and kinematic waves: a study with a numerical ice-flow model. J Glaciol, 41(137), 142-152, https://doi.org/10.3189/S0022143000017834, 1995.

- Van Meijgaard, E., van Ulft, L. H., Lenderink, G., de Roode, S. R., Wipfler, L., Boers, R., and Timmermans, R. M. A.:
 Refinement and application of a regional atmospheric model for climate scenario calculations of Western Europe, No. KVR 054/12, KVR, ISBN EAN 978-90-8815-046-3, 2012.
 Van Vuuren, D. P., Meinshausen, M., Plattner, G. K., Joos, F., Strassmann, K. M., Smith, S. J., Wigley, T. M. L., Raper, S. C. B., Riahi, K., de la Chesnaye, F., den Elzen, M. G. J., Fujino, J., Jiang, K., Nakicenovic, N., Paltsev, S., and Reilly, J. M.: Temperature increase of 21st century mitigation scenarios, P Natl Acad Sci Usa, 105, 15258-15262,
- 475 https://doi.org/10.1073/pnas.0711129105, 2008. World Glacier Monitoring Service (WGMS): Global Glacier Change Bulletin No. 2 (2014-2015). Zemp, M., Nussbaumer, S.U., Gärtner-Roer, I., Huber, J., Machguth, Н., Paul, F. and Hoelzle, (eds.), M. ICSU(WDS)/IUGG(IACS)/UNEP/UNESCO/WMO, World Glacier Monitoring Service, Zurich, Switzerland, 244 pp. Based on database version: https://doi.org/10.5904/wgms-fog-2017-10, 2017 and earlier reports.
- Žebre, M., Colucci, R., Giorgi, F., Glasser, N. F., Racoviteanu, A. E., and Del Gobbo, C.: 200 years of equilibrium-line altitude variability across the European Alps (1901-2100), Clim Dynam, 1-19, https://doi.org/10.1007/s00382-020-05525-7, 2020. Zecchetto, S., Serandrei-Barbero, R., and Donnici, S.: Temperature reconstruction from the length fluctuations of small glaciers in the eastern Alps (northeastern Italy), Clim Dynam, 49(1-2), 363-374, https://doi.org/10.1007/s00382-016-3347-5, 2017.
- 485 Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble, The Cryosphere, 13(4), 1125-1146, https://doi.org/10.5194/tc-13-1125-2019, 2019. Zemp, M., Haeberli, W., Hoelzle, M., and Paul, F.: Alpine glaciers to disappear within decades?, Geophys Res Lett, 33(13), https://doi.org/10.1029/2006GL026319, 2006.
- Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlstroem, A.
 P., Anderson, B., Bajracharya, S., Baroni, C., Brawn, L. N., Caceres, B. E., Casassa, G., Cobos, G., Davila, L. R., Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Haghen, J. O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V., Severskiy, I., Sigurdsson, O., Soruco, A., Usubaliev, R., and Vincent, C.: Historically unprecedented global glacier decline in the early 21st century, J. Glaciol, 61(228), 745-762, 2015.







495

500

Captions

Figure 1: The position of the glaciers and in-situ climatological stations considered in this work. Circles indicate the stations providing air temperature, crosses the precipitation.

Figure 2: Time series of experimental air temperature variations (left panel) and annual total precipitation (right panel) from the in-situ stations available.

Figure 3: The ensemble air temperatures variations over the six regional models at the grid points closest the glaciers position from 2018 to 2100. Left panel: RCP4.5 scenario. Right panel: RCP8.5 scenario. The colours are sorted from the longest (blue) to the shortest (red) glacier.

Figure 4: C_s versus the glaciers slope s for the 22 valley glaciers considered in this work. The solid line is the linear best fit. 505 The glaciers in the legend are sorted from the longest (blue) to the shortest (red).

Figure 5. Observed glacier length variation respect to 1980 length. The colours are sorted from the longest (blue) to the shortest (red) glacier.

Figure 6. The percent glaciers length variations derived from climatological projections from 2018 to 2100. Top panel: RCP4.5 scenario. Bottom panel: RCP8.5 scenario. The colours are sorted from the longest (blue) to the shortest (red) glacier.

510 Figura 7. Ensemble values of σ_L/L derived from climatological projections from 2018 to 2100. Top panel: RCP4.5 scenario. Bottom panel: RCP8.5 scenario. The colours are sorted from the longest (blue) to the shortest (red) glacier. Figure 8: The mean percent glacier length reduction from 2018 to 2100 according to the RCP4.5 and RCP8.5 climatological scenarios.

Table 1. Main parameters of the 22 valley glaciers considered. Slope is the angle between the glacier surface and the horizon.

515 The values of climate sensitivity Cs and response time τ were computed by Eqs. 2 and 3.